





REPLACEMENT SHEET

1/38



ELECTRON-RICH ARYLCHALCOGENO GROUPS WITH DIFFERENT SOLUBILITY CHARACTERISTICS.

FIG. 1A



CORE GROUPS WITH DIFFERENT ATTACHMENT CHARACTERISTICS.

FIG. 1B

BEST AVAILABLE COPY

REPLACEMENT SHEET

2/38




FIG. 2

REPLACEMENT SHEET

3/38




FIG. 3

REPLACEMENT SHEET

4/38




FIG. 4

5/38



REPLACEMENT SHEET

6/38




FIG. 6

REPLACEMENT SHEET

7/38




FIG. 7

8/38



REPLACEMENT SHEET

9/38




FIG. 9

REPLACEMENT SHEET

10/38




FIG. 10


11/38



REPLACEMENT SHEET

12/38

FIG. 12




13/38



REPLACEMENT SHEET

14/38

FIG. 14



REPLACEMENT SHEET

15/38





FIG. 15

REPLACEMENT SHEET

16/38

FIG. 16



REPLACEMENT SHEET

17/38




FIG. 17





FIG. 18

REPLACEMENT SHEET

18/38

FIG. 19



REPLACEMENT SHEET

19/38




FIG. 20

REPLACEMENT SHEET

20/38




FIG. 21A



FIG. 21B

REPLACEMENT
SHEET

21/38



FIG. 21C



FIG. 21D

REPLACEMENT SHEET

22/38




FIG. 21E



FIG. 21F

REPLACEMENT SHEET

23/38




FIG. 22A




FIG. 22B

REPLACEMENT SHEET

24/38

DISTRIBUTION OF ENTEROMORPHA ZOOSPORES AFTER
EXPOSURE TO A SURFACE PRESSURE OF 56 KPa




FIG. 23

PERCENTAGE REMOVAL OF ENTEROMORPHA ZOOSPORES AFTER
EXPOSURE TO A SURFACE PRESSURE OF 56 KPa




FIG. 24A

REPLACEMENT SHEET

25/38




FIG. 24B




FIG. 25

REPLACEMENT SHEET

26/38




FIG. 26

REPLACEMENT SHEET

27/38




FIG. 27




FIG. 28

REPLACEMENT SHEET

28/38




FIG. 29




FIG. 30

REPLACEMENT SHEET

29/38




FIG. 31




FIG. 32

REPLACEMENT SHEET

30/38




FIG. 33




FIG. 34

REPLACEMENT SHEET

31/38

DENSITY OF ENTEROMORPHA SPORES REMAINING AFTER
PRESSURE OF 83 kPa UNDER WATER JET




FIG. 35

PERCENT REMOVAL OF ENTEROMORPHA SPORES FROM SOL-GELS
AFTER PRESSURE OF 83 kPa UNDER WATER JET




FIG. 36

REPLACEMENT SHEET

32/38




FIG. 37




FIG. 38

REPLACEMENT SHEET

33/38




FIG. 39




FIG. 40

REPLACEMENT SHEET

34/38




FIG. 41

REPLACEMENT SHEET

35/38




FIG. 42




FIG. 43

REPLACEMENT SHEET

36/38




FIG. 44




FIG. 45

REPLACEMENT
SHEET

37/38




FIG. 46

REPLACEMENT SHEET

38/38




FIG. 47




FIG. 48

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ ~~SKewed/SLANTED IMAGES~~
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.